optionally substituted heteroaryl-, optionally substituted heterocyclyl-, acyl-, alkoxycarbonyl-, sulfanyl-, sulfinyl and sulfonyl-, e.g., diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino.

[0084] Aminocarbonyl- refers to the group —NR°COR^b, —NR°CO₂R^a, or —NR°CONR^bR°, where

[0085] R^1 is optionally substituted C_1 - C_6 alkyl-, aryl-, heteroaryl-, aryl- C_1 - C_4 alkyl-, or heteroaryl- C_1 - C_4 alkyl-group;

[0086] R^b is H or optionally substituted C_1 - C_6 alkyl-, aryl-, heteroaryl-, aryl- C_1 - C_4 alkyl-, or heteroaryl- C_1 - C_4 alkyl- group; and

[0087] R° is hydrogen, alkyl-; aryl- or heteroaryl-; and where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C₁-C₄ alkyl-, aryl-, heterocyclyl-, aryl- C_1 - C_4 alkyl-, heteroaryl- C_1 - C_4 alkyl-, C_1 - C_4 haloalkyl-, $-OC_1$ - C_4 alkyl, $-OC_1$ - C_4 alkylphenyl, $-C_1$ -C₄ alkyl-OH, —OC₁-C₄ haloalkyl, halogen, —OH, —NH₂, —NR°C(NRb)(NRbR°) (i.e. guanidine), —NR°CRbNRbR°, $-\text{CNR}^{c}\text{NR}^{b}\text{R}^{c}$, $-\text{C}_{1}\text{-C}_{4}$ alkyl-NH₂, $-\text{N}(\text{C}_{1}\text{-C}_{4}$ alkyl)(C₁- C_4 alkyl), —NH(C_1 - C_4 alkyl), —N(C_1 - C_4 alkyl)(C_1 - C_4 alkylphenyl), —NH(C₁-C₄ alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), — CO_2H , — $C(O)OC_1-C_4$ alkyl-, — $CON(C_1-C_4$ alkyl)(C_1-C_4 alkyl), — $CONH(C_1-C_4$ alkyl), —CONH₂, —NHC(O)(C₁-C₄ alkyl), —NH-C(O)(phenyl), $-N(C_1-C_4 \text{ alkyl})C(O)(C_1-C_4 \text{ alkyl})$, $-N(C_1-C_4 \text{ alkyl})$ C_4 alkyl)C(O)(phenyl), —C(O) C_1 - C_4 alkyl-, —C(O) C_1 - C_4 phenyl-, $-C(O)C_1-C_4$ haloalkyl-, $-OC(O)C_1-C_4$ alkyl-, $-SO_3(C_1-C_4)$ alkyl), —SO₂(phenyl), $-SO_2(C_1-C_4)$ $-SO_2NH_2$, $-SO_2NH(C_1-C_4)$ alkyl), haloalkyl), -NHSO₂(C₁-C₄ —SO₂NH(phenyl), alkyl), —NHSO₂(phenyl), and —NHSO₂(C₁-C₄ haloalkyl).

[0088] Antimitotic refers to a drug for inhibiting or preventing mitosis, for example, by causing metaphase arrest. Some antitumour drugs block proliferation and are considered antimitotics.

[0089] Aryl and heteroaryl mean a 5- or 6-membered aromatic or heteroaromatic ring containing 0 or 1-4 heteroatoms, respectively, selected from O, N, or S; a bicyclic 9- or 10-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms, respectively, selected from O, N, or S; or a tricyclic 12- to 14-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms, respectively, selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., phenyl-, naphthyl-, indanyl-, tetralinyl-, and fluorenyl and the 5- to 10-membered aromatic heterocyclic rings include, e.g., imidazolyl-, pyridinyl-, indolyl-, thienyl-, benzopyranonyl-, thiazolyl-, furanyl-, benzimidazolyl-, quinolinyl-, isoquinolinyl-, quinoxalinyl-, pyrimidinyl-, pyrazinyl-, tetrazolyl and pyrazolyl-.

[0090] Aralkyl- refers to a residue in which an aryl moiety is attached to the parent structure via an alkyl residue. Examples include benzyl-, phenethyl-, phenylvinyl-, phenylallyl and the like. Heteroaralkyl- refers to a residue in which a heteroaryl moiety is attached to the parent structure via an alkyl residue. Examples include furanylmethyl-, pyridinylmethyl-, pyrimidinylethyl and the like.

[0091] Aralkoxy- refers to the group —O-aralkyl. Similarly, heteroaralkoxy- refers to the group —O-het-

eroaralkyl-; aryloxy- refers to the group —O-aryl-; acyloxy-refers to the group —O-acyl-; heteroaryloxy- refers to the group —O-heteroaryl-; and heterocyclyloxy- refers to the group —O-heterocyclyl (i.e., aralkyl-, heteroaralkyl-, aryl-, acyl-, heterocyclyl-, or heteroaryl is attached to the parent structure through an oxygen).

[0092] Carboxyalkyl- refers to the group -alkyl-COOH.

[0093] Carboxamido refers to the group —CONR $^{b}R^{c},$ where

[0094] R^b is H or optionally substituted C_1 - C_6 alkyl-, aryl-, heteroaryl-, aryl- C_1 - C_4 alkyl-, or heteroaryl- C_1 - C_4 alkyl- group; and

[0095] R° is hydrogen, alkyl-; aryl- or heteroaryl-; and

[0096] where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C₁-C₄ alkyl-, aryl-, heterocyclyl-, aryl-C₁-C₄ alkyl-, heteroaryl-C₁-C₄ alkyl-, C_1 - C_4 haloalkyl-, — OC_1 - C_4 alkyl-, — OC_1 - C_4 alkylphenyl-, $^{\perp}C_1$ - C_4 alkyl-OH, —OC₁- C_4 haloalkyl-, halogen, —OH, $\begin{array}{l} -\mathrm{NH_2}, -\mathrm{C_1\text{-}C_4} \text{ alkyl-NH_2}, -\mathrm{N(C_1\text{-}C_4} \text{ alkyl)} (\mathrm{C_1\text{-}C_4} \text{ alkyl)}, \\ -\mathrm{NH(C_1\text{-}C_4} \text{ alkyl)}, -\mathrm{N(C_1\text{-}C_4} \text{ alkyl)} (\mathrm{C_1\text{-}C_4} \text{ alkyl)} (\mathrm{C_1\text{-}C_4} \text{ alkyl)}, \end{array}$ -NH(C₁-C₄ alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), $-CO_2H$, $-C(0)OC_1-C_4$ alkyl-, $-CON(C_1-C_4)$ alkyl), $-CONH(C_1-C_4)$ alkyl), $-CONH(C_1-C_4)$ alkyl), $-CONH(C_1-C_4)$ $-NHC(O)(C_1-C_4 \text{ alkyl}), -NHC(O)(\text{phenyl}), -N(C_1-C_4)$ alkyl) $C(O)(C_1^1-C_4$ alkyl), $-N(C_1-C_4$ alkyl)C(O)(phenyl), $-C(O)C_1-C_4$ alkyl-, $-C(O)C_1-C_4$ phenyl-, $-C(O)C_1-C_4$ haloalkyl-, $-OC(O)C_1$ - C_4 alkyl-, $-SO_2(C_1$ - C_4 alkyl), $-SO_2(\text{phenyl})$, $-SO_2(C_1$ - C_4 haloalkyl), $-SO_2NH_2$, $-SO_2NH(C_1$ - C_4 haloalkyl), $-SO_2NH(C_1$ - C_4 haloalkyl), $-SO_2NH(C_1$ - C_4 haloalkyl), $-SO_2NH(C_1$ - $-SO_2NH(C_1$ -- C_4 alkyl), —NHSO₂(phenyl), and —NHSO₂(\tilde{C}_1 - C_4 haloalkyl). Carboxamido is meant to include carbamoyl-; lower-alkyl carbamoyl-; benzylcarbamoyl-; phenylcarbamoyl-; methoxymethyl-carbamoyl-; and the like.

[0097] Halogen or halo refers to fluorine, chlorine, bromine or iodine. Fluorine, chlorine and bromine are preferred. Dihaloaryl-, dihaloalkyl-, trihaloaryl etc. refer to aryl and alkyl substituted with the designated plurality of halogens (here, 2, 2 and 3, respectively), but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl-.

[0098] Heterocyclyl means a cycloalkyl or aryl residue in which one to foul of the carbons is replaced by a heteroatom such as oxygen, nitrogen or sulfur. Examples of heterocycles that fall within the scope of the invention include azetidinyl-, imidazolinyl-, pyrrolidinyl-, pyrazolyl-, pyrrolyl-, indolyl-, quinolinyl-, isoquinolinyl-, tetrahydroisoquinolinyl-, benzofuranyl-, benzodioxanyl-, benzodioxyl (commonly referred to as methylenedioxyphenyl-, when occurring as a substituent), tetrazolyl-, morpholinyl-, thiazolyl-, pyridinyl-, pyridazinyl-, piperidinyl-, pyrimidinyl-, thienyl-, furanyl-, oxazolyl-, oxazolinyl-, isoxazolyl-, dioxanyl-, tetrahydrofuranyl and the like. "N-heterocyclyl" refers to a nitrogencontaining heterocycle. The term heterocyclyl encompasses heteroaryl-, which is a subset of heterocyclyl-. Examples of N-heterocyclyl residues include azetidinyl-, 4-morpholinyl-, 4-thiomorpholinyl-, 1-piperidinyl-, 1-pyrrolidinyl-, 3-thiazolidinyl-, piperazinyl and 4-(3,4-dihydrobenzoxazinyl). Examples of substituted heterocyclyl include 4-methyl-1piperazinyl and 4-benzyl-1-piperidinyl-.